

Explaining link between emotion and addictive substance use

Date:

January 9, 2020

Source:

Harvard Kennedy School

Summary:

What drives a person to smoke cigarettes? What role do emotions play in this addictive behavior? Why do some smokers puff more often and more deeply or relapse many years after they've quit? If policy makers had those answers, how could they strengthen the fight against the global smoking epidemic? A new report by researchers offers a key insight: sadness plays an especially strong role in triggering addictive behavior relative to other negative emotions like disgust.

FULL STORY

What drives a person to smoke cigarettes -- and keeps one out of six U.S. adults addicted to tobacco use, at a cost of 480,000 premature deaths each year despite decades of anti-smoking campaigns? What role do emotions play in this addictive behavior? Why do some smokers puff more often and more deeply or even relapse many years after they've quit? If policy makers had those answers, how could they strengthen the fight against the global smoking epidemic?

advertisement

A team of researchers based at Harvard University now has fresh insights into these questions, thanks to a set of four interwoven studies described in a new report published in the *Proceedings of the National Academy of Sciences*: The studies show that sadness plays an especially strong role in triggering addictive behavior relative to other negative emotions like disgust.

The studies range from analysis of data from a national survey of more than 10,000 people over 20 years to laboratory tests examining the responses of current smokers to negative emotions. One study tested the volume and frequency of actual puffs on cigarettes by smokers who volunteered to be monitored as they smoked. While drawing from methodologies from different

fields, the four studies all reinforce the central finding that sadness, more so than other negative emotions, increases people's craving to smoke.

"The conventional wisdom in the field was that any type of negative feelings, whether it's anger, disgust, stress, sadness, fear, or shame, would make individuals more likely to use an addictive drug," said lead researcher Charles A. Dorison, a Harvard Kennedy School doctoral candidate. "Our work suggests that the reality is much more nuanced than the idea of 'feel bad, smoke more.' Specifically, we find that sadness appears to be an especially potent trigger of addictive substance use."

Senior co-author Dr. Jennifer Lerner, the co-founder of the Harvard Decision Science Laboratory and Thornton F. Bradshaw Professor of Public Policy, Decision Science, and Management at Harvard Kennedy School, said the research could have useful public policy implications. For example, current anti-smoking ad campaigns could be redesigned to avoid images that trigger sadness and thus unintentionally increase cigarette cravings among smokers.

Lerner is the first tenured psychologist on the faculty of the Kennedy School. She was the Chief Decision Scientist for the U.S. Navy in 2018-19. Lerner has studied the impact of emotions on decision-making since the 1990s, examining issues including whether generalized negative emotions trigger substance abuse or whether a subset of specific emotions such as sadness are more important factors in addiction.

The other co-authors include Ke Wang, a doctoral student at the Kennedy School; Vaughan W. Rees, director of the Center for Global Tobacco Control at Harvard T.H. Chan School of Public Health; Ichiro Kawachi, the John L. Loeb and Frances Lehman Loeb Professor of Social Epidemiology at the Chan School; and Associate Professor Keith M.M. Ericson at the Questrom School of Business at Boston University. The work was funded by grants from the National Science Foundation and the National Institutes of Health.

Here are further details on the techniques and key findings of the four studies:

1. Examining data from a national survey that tracked 10,685 people over 20 years, the researchers found that self-reported sadness among participants was associated with being a smoker and with relapsing back into smoking one and two decades later. The sadder individuals were, the more likely they were to be smokers. Notably, other negative emotions did not show the same relationship with smoking.
2. Then the team designed an experiment to test causality: Did sadness cause people to smoke, or were negative life events causing both sadness and smoking? To test this, 425 smokers were recruited for an online study: one-third were shown a sad video clip about the loss of a life partner. Another third of the smokers were shown a neutral video clip, about woodworking; the final third were shown a disgusting video involving an unsanitary toilet. All participants were asked to write about a related personal experience. The study found that individuals in the sadness condition -- who watched the sad video and wrote about a personal loss -- had higher cravings to smoke than both the neutral group and the disgust group.
3. A similar approach in the third study measured actual impatience for cigarette puffs rather than mere self-reported craving. Similar to the second study, nearly 700

participants watched videos and wrote about life experiences that were either sad or neutral, and then were given hypothetical choices between having fewer puffs sooner or more puffs after a delay. Those in the sadness group proved to be more impatient to smoke sooner than those in the neutral group. That result built upon previous research findings that sadness increases financial impatience, measured with behavioral economics techniques.

4. The fourth study recruited 158 smokers from the Boston area to test how sadness influenced actual smoking behavior. Participants had to abstain from smoking for at least eight hours (verified by carbon monoxide breath test). They were randomly assigned to sadness or neutral control groups; smokers sat in a private room at the Harvard Tobacco Research Laboratory, watched the sad video and wrote about great loss, or watched a neutral video and wrote about their work environment. Then they smoked their own brand through a device that tested the total volume of puffs and their speed and duration. The results: smokers in the sadness condition made more impatient choices and smoked greater volumes per puff.

Lerner said the research team was motivated in part by the deadly realities of smoking: tobacco use remains the leading cause of preventable death in the United States despite five decades of anti-smoking campaigns. The global consequences are also dire, with one billion premature deaths predicted across the world by the end of this century.

"We believe that theory-driven research could help shed light on how to address this epidemic," Dorison said. "We need insights across disciplines, including psychology, behavioral economics and public health, to confront this threat effectively."

Story Source:

[Materials](#) provided by [Harvard Kennedy School](#). Original written by James F. Smith. *Note: Content may be edited for style and length.*

Journal Reference:

1. Charles A. Dorison, Ke Wang, Vaughan W. Rees, Ichiro Kawachi, Keith M. M. Ericson, Jennifer S. Lerner. **Sadness, but not all negative emotions, heightens addictive substance use.** *Proceedings of the National Academy of Sciences*, 2019; 201909888 DOI: [10.1073/PNAS.1909888116](https://doi.org/10.1073/PNAS.1909888116)

Cite This Page:

- [MLA](#)
- [APA](#)
- [Chicago](#)

Harvard Kennedy School. "Explaining link between emotion and addictive substance use." ScienceDaily. ScienceDaily, 9 January 2020.

<www.sciencedaily.com/releases/2020/01/200109152823.htm>.

Harvard Kennedy School. (2020, January 9). Explaining link between emotion and addictive substance use. *ScienceDaily*. Retrieved January 10, 2020 from www.sciencedaily.com/releases/2020/01/200109152823.htm

Harvard Kennedy School. "Explaining link between emotion and addictive substance use." ScienceDaily. www.sciencedaily.com/releases/2020/01/200109152823.htm (accessed January 10, 2020).

RELATED STORIES

[Blunting Pain's Emotional Component](#)

Mar. 13, 2019 — Pain researchers have shown in rodents that they can block receptors on brain cells that appear to be responsible for the negative emotions associated with pain, such as sadness, depression and ... [read more](#)

[Marijuana May Lead Non-Smokers to Cigarettes](#)

Mar. 27, 2018 — While cigarette smoking is declining, marijuana use is rising and, disproportionately, marijuana users also smoke cigarettes. Our new study reports that cannabis use was associated with an increased ... [read more](#)

[A Mom's Support Helps a Child Learn to Handle Negative Emotions, but What If Mom Is Distressed?](#)

Nov. 17, 2017 — When children become upset, showing negative emotions or behaviors, some parents become distressed, while others are able to talk their child through the difficult situation. Studies have shown that ... [read more](#)

[Managing Negative Emotions Can Help Pregnant Smokers Quit](#)

Sep. 19, 2017 — Pregnant smokers are more likely to quit if they can learn to manage negative emotions that lead to smoking, new research ... [read more](#)